

Lecture Notes in Mathematics

Edited by A. Dold and B. Eckmann

1364

Robert R. Phelps

Convex Functions,
Monotone Operators
and Differentiability



Springer-Verlag Berlin Heidelberg GmbH

Author

Robert R. Phelps
Department of Mathematics GN-50, University of Washington
Seattle, WA 98195, USA

Mathematics Subject Classification (1980): 46B20, 46B22, 47H05, 49A29,
49A51, 52A07

ISBN 978-3-540-50735-2 ISBN 978-3-662-21569-2 (eBook)
DOI 10.1007/978-3-662-21569-2

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, re-use of illustrations, recitation, broadcasting, reproduction on microfilms or in other ways, and storage in data banks. Duplication of this publication or parts thereof is only permitted under the provisions of the German Copyright Law of September 9, 1965, in its version of June 24, 1985, and a copyright fee must always be paid. Violations fall under the prosecution act of the German Copyright Law.

© Springer-Verlag Berlin Heidelberg 1989

Originally published by Springer-Verlag Berlin Heidelberg New York in 1989

2146/3140-543210

PREFACE

These notes had their genesis in a widely distributed but unpublished set of notes *Differentiability of convex functions on Banach spaces* which I wrote in 1977-78 for a graduate course at University College London (UCL). Those notes were largely incorporated into J. Giles' 1982 Pitman Lecture Notes *Convex analysis with application to differentiation of convex functions*. In the course of doing so, he reorganized the material somewhat and took advantage of any simpler proofs available at that time. I have not hesitated to return the compliment by using a few of those improvements. At my invitation, R. Bourgin has also incorporated material from the UCL notes in his extremely comprehensive 1983 Springer Lecture Notes *Geometric aspects of convex sets with the Radon-Nikodym property*. The present notes do not overlap too greatly with theirs, partly because of a substantially changed emphasis and partly because I am able to use results or proofs that have come to light since 1983.

Except for some subsequent revisions and modest additions, this material was covered in a graduate course at the University of Washington in Winter Quarter of 1988. The students in my class all had a good background in functional analysis, but there is not a great deal needed to read these notes, since they are largely self-contained; in particular, no background in convex functions is required. The main tool is the separation theorem (a.k.a. the Hahn-Banach theorem); like the standard advice given in mountaineering classes (concerning the all-important bowline for tying oneself into the end of the climbing rope), you should be able to employ it using only one hand while standing blindfolded in a cold shower.

These notes have been influenced very considerably by frequent conversations with Isaac Namioka (who has an almost notorious instinct for simplifying proofs) as well as occasional conversations with Terry Rockafellar; I am grateful to them both. I am also grateful to Jon Borwein, Marian Fabian and Simon Fitzpatrick, each of whom sent me useful suggestions based on a preliminary version.

Robert R. Phelps
October 5, 1988
Seattle, Washington

Production note: I typed these notes on a Macintosh using MacWrite 4.5 (with the *Princeton* Font 2.0) for the text and MacPaint for the drawings. The non-mathematical portions (such as the present page) were done in the *New York* font and all of it was printed on an Apple LaserWriter II.

INTRODUCTION

The study of the differentiability properties of convex functions on infinite dimensional spaces has continued on and off for over fifty years. There are a couple of obvious reasons for this. Aside from the intrinsic interest of investigating the many consequences implicit in something as simple as convexity, there is the satisfaction (for this author, at least) in discovering that a number of apparently disparate mathematical topics (extreme points - rather, strongly exposed points - of noncompact convex sets, monotone operators, perturbed optimization of real-valued functions, differentiability of vector-valued measures) are in fact closely intertwined, with differentiability of convex functions forming a common thread.

Starting in Section 1 with the definition of convex functions and a fundamental differentiability property in the one-dimensional case [right-hand and left-hand derivatives always exist], we get quickly to the first infinite dimensional result, Mazur's intriguing 1933 theorem: A continuous convex function on a separable Banach space has a dense G_δ set of points where it is (Gateaux) differentiable. In order to go beyond Mazur's theorem, some time is spent in studying the subdifferential of a convex function f ; this is a set-valued map from the space to its dual whose image at each point x consists of all plausible candidates for the derivative of f at x . [The function f is Gateaux differentiable precisely when the subdifferential is single-valued, and it is Fréchet differentiable precisely when its subdifferential is single-valued and norm-to-norm continuous.]

Since a subdifferential is a special case of a monotone operator, Section 2 starts with a detailed look at monotone operators. These objects are of independent origin, having been extensively studied in the sixties and early seventies by numerous mathematicians (with major contributions from H. Brezis, F. Browder and G. J. Minty) in connection with nonlinear partial differential equations and other aspects of nonlinear analysis. (See, for instance, [Bre] or [Pa-Sb]). Also in the sixties, an in-depth study of monotone operators in fairly general spaces was carried out by R. T. Rockafellar, who established a number of fundamental properties, such as their local boundedness. He also gave an elegant characterization of those monotone operators which are the subdifferentials of convex functions, a theorem which is much easier to state than to prove (and which is not proved in full generality until Section 3). [The connection between monotone operators and derivatives of convex functions is readily apparent on the real line, since monotone operators coincide in that case with monotone nondecreasing functions, as do the right-hand derivatives of convex functions of one variable.]

In 1968, E. Asplund extended Mazur's theorem in two ways: He found more general spaces in which the same conclusion holds, and he studied a less general class of Banach spaces (now called Asplund spaces) in which a stronger conclusion holds. (Namely, he replaced the Gateaux derivative by the stronger Fréchet derivative.) Asplund used an ingenious combination of analytic and geometric techniques to prove some of the basic theorems in the subject. Roughly ten years later, P. Kenderov (as well as R. Robert and S. Fitzpatrick) proved some general continuity theorems for monotone operators which, when applied to subdifferentials, yield Asplund's results as special cases. In Section 2 we follow this approach, incorporating recent work by D. Preiss and L. Zajicek to obtain the major differentiability theorems.

The results of Section 2 all involve continuous convex functions defined on open convex sets. For many applications, it is more suitable to consider lower semicontinuous convex functions, even those which are extended real valued (possibly equal to $+\infty$). (For instance, in many optimization problems one finds just such a function in the form of the supremum of an infinite family of affine continuous functions.) Lower semicontinuous convex functions also yield a natural way to translate results about closed convex sets into results about convex functions and vice versa. (For instance, the set of points on or above the graph of such a convex function - its epigraph - forms a closed convex set). In Section 3 one will find some classical results (various versions and extensions of the Bishop-Phelps theorems) which, among other things, guarantee that subdifferentials still exist for lower semicontinuous convex functions. A nonconvex version of this type of theorem is I. Ekeland's variational principle, which asserts that a lower semicontinuous function which nearly attains its minimum at a point x admits arbitrarily small perturbations (by translates of the norm) which do attain a minimum, at points near x . This result, while simple to state and prove, has been shown by Ekeland [Ek] to have an extraordinarily wide variety of applications, in areas such as optimization, mathematical programming, control theory, nonlinear semigroups and global analysis.

In Section 4, a variational principle is established which uses differentiable perturbations; this recent result is due to J. Borwein and D. Preiss. Some deep theorems about differentiability of convex functions fall out as fairly easy corollaries, and it is reasonable to expect future useful applications.

Section 5 describes the duality between Asplund spaces and spaces with the Radon-Nikodym property (RNP). These are Banach spaces for which a Radon-Nikodym-type differentiation theorem is valid for vector measures with values in the space. Spaces with the RNP have an interesting history, starting in the late sixties with the introduction by M. Rieffel of a geometric property (dentability) which turned out to characterize the RNP and which

has led to a number of other characterizations in terms of the extreme points (or strongly exposed points) of bounded closed, convex subsets of the space. A truly beautiful result in this area is the fact that a Banach space is an Asplund space if and only if its dual has the RNP. (Superb expositions of the RNP may be found in the books by J. Diestel and J. J. Uhl [Di-U] and R. Bourgin [Bou].) In Section 5, the RNP is defined in terms of dentability, and a number of basic results are obtained using more recent (and simpler) proofs than are used in the above monographs. One will also find there J. Bourgain's proof of C. Stegall's perturbed optimization theorem for semicontinuous functions on spaces with the RNP; this yields as a corollary the theorem that in such spaces every bounded closed convex set is the closed convex hull of its strongly exposed points.

The notion of perturbed optimization has been moving closer to center stage, since it not only provides a more general format for stating previously known theorems, but also permits the formulation of more general results. The idea is simple: One starts with a real-valued function f which is, say, lower semicontinuous and bounded below on a nice set, and shows that there exist arbitrarily small perturbations g such that $f + g$ attains a minimum on the set. The perturbations g might be restrictions of continuous linear functionals of small norm, or perhaps Lipschitz functions of small Lipschitz norm. Moreover, for really nice sets, the perturbed function attains a strong minimum: Every minimizing sequence converges.

The brief Section 6 is devoted to the class of Banach spaces in which every continuous convex function is Gateaux differentiable in a dense set of points (dropping the previous condition that the set need be a G_δ). Some evidence is presented that this is perhaps the "right" class to study.

Even more general than monotone operators is a class of set valued maps (from a metric space, say, to a dual Banach space) which are upper semicontinuous and take on weak* compact convex values, the so-called usco maps. In Section 7, some interesting connections between monotone operators and usco maps are described, culminating in a topological proof of one of P. Kenderov's continuity theorems.

CONTENTS

| | |
|--|------------|
| 1. Convex functions on real Banach spaces | 1 |
| Subdifferentials of continuous convex functions, Gateaux and Fréchet differentiability, Mazur's theorem. | |
| 2. Monotone operators, subdifferentials and Asplund spaces | 17 |
| Upper semicontinuity of set-valued monotone operators, characterization of Gateaux and Fréchet differentiability in terms of continuous selections, Preiss-Zajicek generic continuity theorem for monotone operators into separable dual spaces, Asplund spaces and subspaces with separable duals, weak*-slices, subdifferentials of convex functions as maximal monotone operators, local boundedness of monotone operators, Kenderov's generic continuity theorems for maximal monotone operators, weakly compactly generated dual spaces and Asplund spaces. | |
| 3. Lower semicontinuous convex functions | 40 |
| Extended real-valued convex functions and their subdifferentials, support points, minimal points with respect to special cones, Ekeland's variational principle, Borwein's unification, Brøndsted-Rockafellar theorem, Bishop-Phelps theorems, maximal monotonicity of the subdifferential, maximal cyclically monotone operators are subdifferentials, subdifferentials of saddle functions. | |
| 4. A smooth variational principle and more about Asplund spaces | 64 |
| Borwein-Preiss smooth variational principle, Ekeland-Lebourg theorem. | |
| 5. Asplund spaces, the Radon-Nikodym property and optimization | 72 |
| Slices and weak* slices and dentability, RNP, infinite trees, E is an Asplund space means E^* has the RNP, duality between weak* strongly exposed points and Fréchet differentiability, perturbed optimization on RNP sets, bounded closed convex RNP sets are generated by their strongly exposed points, Ghoussoub-Maurey theorems. | |
| 6. Gateaux differentiability spaces. | 90 |
| Gateaux differentiability spaces, equivalence with M -differentiability spaces, duality characterization in terms of weak* exposed points, stability results | |
| 7. A generalization of monotone operators: Usco maps | 97 |
| Upper semicontinuous compact valued (usco) maps, maximal monotone operators are minimal usco maps, topological proof of Kenderov's generic continuity theorem. | |
| 8. Notes and Remarks | 104 |
| Background, section by section, as well as suggestions for further reading. | |
| References | 108 |
| Index | 113 |
| Index of Symbols | 115 |